

# Special Research Report # 106: Disease Management

## Biological Control of *Rhizoctonia* Root Rot

D.J. Norman, Associate Professor, Department of Plant Pathology  
University of Florida, MREC-Apopka, FL 32779



FUNDING INDUSTRY SOLUTIONS  
TODAY & TOMORROW

Phone: 618/692-0045

Fax: 618/692-4045

E-mail: [afe@endowment.org](mailto:afe@endowment.org)

Website: [www.endowment.org](http://www.endowment.org)

### BACKGROUND

Most flowering plants are propagated via seeds or cuttings that are established in seeding production trays that contain from 72 to over 500 individual plug units per tray. Each unit is filled with an artificial planting media usually consisting of a mixture of peatmoss, perlite,



Dave J. Norman

[DJN@gvn.ifas.ufl.edu](mailto:DJN@gvn.ifas.ufl.edu)

407-884-2034 ext 150

and/or vermiculite. Even with the use of this artificial

media, extensive production losses can be caused by *Rhizoctonia* root rot.

This pathogen appears to gain access to facilities via contaminated seed, infested cuttings, or unsanitary growing practices.

The objectives of this research were to: (1) Identify *Bacillus* spp. that naturally colonize roots of bedding plants; (2) Identify which of these *Bacillus* spp. might be useful for biological control of *Rhizoctonia* or for plant growth promotion

### MATERIALS AND METHODS

**Isolation.** Root samples were collected from 35 genera of annual and perennial bedding plants in well established planting beds. *Bacillus* spp. were selectively isolated using heat treatment and a selective medium. A total of 149 *Bacillus* strains were identified to species using fatty acid analyses.

**Evaluation.** All 149 *Bacillus* strains were screened with the following procedures. *Bacillus* strains were cultured, quantified, and mixed into a medium

consisting of 60% Canadian peat moss, 20% vermiculite and 20% perlite.

Impatiens plants are the most common bedding plants grown in the US and were selected for this study because of the following: (1) Could easily be used in mechanical seeding; (2) the germination of impatiens seeds (4 days) and growth of seedlings are rapid; and (3) plants exhibit complete damping-off when infected with *Rhizoctonia* allowing for ease of evaluation.



Seedling damping-off


Impatiens seeds ('Super Elfin White') were seeded into 288 cell plug trays. Each tray contained the planting mix



Seedlings being grown with *Bacillus* isolates

and one of the 149 *Bacillus* isolates. Seeds were grown in trays for four weeks and subsequently transplanted into 72 plug seeding trays containing an average of 65 *Rhizoctonia* sclerotia per plug (sclerotia are survival structures formed by *Rhizoctonia*). Number of seedlings damping-off in each treatment were compared on a daily basis for two weeks. Two controls were utilized each time *Bacillus* strains were tested; 1) "Disease control" *Rhizoctonia* – no *Bacillus*, 2) "Fungicide Control" *Rhizoctonia* + Banrot 50WP without *Bacillus*.


Seven strains from seven *Bacillus* spp. were selected from the initial screening. Tests were repeated with these strains a minimum of three times. Means of these *Bacillus* strains were all lower than disease controls (Graph 1).



Graph 1


Three of the seven *Bacillus* strains were selected to test for a synergist effect by applying combinations of two strains to impatiens transplants. All nine

permutations were tested five times. No other changes were made to the described protocol. Combinations of *B. cereus*, *B. lacterosporus*, and *B. pumilis* were synergistic in their suppression of *Rhizoctonia* (Graph 2).



Graph 2

In order to determine if the *Bacillus* strains are promoting plant growth in soilless media; the seven previously selected *Bacillus* strains were incorporated into 100 cell seeding trays as previously described. After four weeks of growth above ground portions of impatiens plants were removed dried and compared with that of a control tray in which no *Bacillus* strain was added. This was repeated three times and data were compared.



Graph 3

## CONCLUSIONS

The fatty acid analysis (MIDI) classified most of the 149 *Bacillus* strains into 14 spp. The seven most common species associated with the roots were: *B. lacterosporus* (24%), *B. cereus* (21%), *B. megaterium* (13%), *B. mycoides* (5%), *B. pumilis* (5%), *B. thuringiensis* (4%), *B. coagulans* (4%). Ability to suppress *Rhizoctonia* by individual strains within these species varied greatly. However, some *Bacillus* strains were found that slowed progression of root rot and promoted growth of the plant.

## IMPACT TO THE INDUSTRY

(1). None of the 149 *Bacillus* strains are capable of suppressing *Rhizoctonia* to the same degree as commercial fungicides; however, in the bedding plant industry, these strains have a potential benefit if applications are made at seeding and transplanting. (2) These strains appear to aid in plant growth when an artificial soilless media is used.

December 2000 © Copyright The American Floral Endowment. All Rights Reserved.